

Chlamydia/Gonorrhea

Two of the most common bacterial infections in Utah

Treatment: Antibiotics

Get the Facts: Utah Statistics

- One in 4 teens infected with Chlamydia during 2012
- Chlamydia/Gonorrhea most commonly found in people ages 15-29

Utah AIDS Foundation

Transmission

- Unprotected vaginal, oral or anal sex
- Passed from mother to baby during birth
- Genital to genital contact (such as mutual masturbation)

Symptoms (may appear 1-3 weeks from exposure)

Women: 75% don't experience symptoms

- Unusual discharge (yellow or blood)
- Itching, burning, and/or pain while peeing
- Bleeding during sex
- Change in period
- Stomach pain/fever

Men: 50% don't experience symptoms

- Discharge from Penis (yellow/green)
- Burning/pain while peeing
- Itching around opening of Penis
- Swollen testicles

Rectal: Localized to this area

- Rectal discharge/bleeding
- Pain

Oral: Localized to this area

- Sore throat/swollen lymph nodes

Tests & Costs

Chlamydia & Gonorrhea are always tested together. Because the bacteria will only infect the area it has come into contact with, there are 3 different tests offered. These include:

Urine Sample \$25	Used to detect bacteria infecting the penis or Vagina
Oral throat swab \$25	Self-collected throat swab to detect bacteria in mouth/throat
Rectal swab \$25	Self-collected rectal swab to detect bacteria in rectum

**KNOW
YOUR
STATUS**

After the test:

Once you have been tested, you will be given a card with a unique ID number that will be used to identify you when you call UAF 2 weeks later to get your results. UAF will only contact you if your results are positive so it is important to call to receive test results.